



**Climate smart inputs for
precision farming!**




APPLICATION SCHEDULE - MAIZE / CORN



Contact us -

 +91 7991 7991 30, +91 9850 9214 48

 +91 9420 3117 76  sales@aryabiotech.com

 www.aryabiotech.com



Application dosage :

Seed treatment - 10 ml per kg of seed

Foliar spray - 3 to 5 ml per liter of water

Soil application - 10 to 25 kg per acre

Stage	Product	Application method
At the time of sowing	NANOWISE STARTER	Seed treatment
	BIOBOOST Powder	Soil application
15 days after sowing	BALADA Liquid	Foliar spray
At the time of flowering	ENGIMINS MINTY	Foliar spray
	BALADA Liquid	Foliar spray
At the time of gram setting	ENGIMINS MINTY	Drip / Drenching

- For protection from pests/diseases apply MAHANIMBA 3 to 5 ml per liter of water by foliar spray.
- ENGIMINS SHIELD should be applied at the time of fungal attack or humid weather, 3 ml per liter of water by foliar spray and 1 liter per acre through drip.
- Package of agri-practices may vary according to geographical, climatic conditions, soil type etc.
- Base your application of ENGIMINS on leaf, petiole or tissue analysis. The lower rates should be used for maintenance while the higher rates will correct nutritional deficiencies.

